

High Resolution Spectroscopy of the $^{12}_{\Lambda}\text{B}$ Hypernucleus Produced by the $(e,e'K^+)$ Reaction

T. Miyoshi^a, M. Sarsour^b, L. Yuan^c, X. Zhu^c, A. Ahmidouch^d, P. Ambrozewicz^e, D. Androic^f, T. Angelescu^g, R. Asaturyan^h, S. Avery^c, O.K. Baker^{c,j}, I. Bertovic^f, H. Breuerⁱ, R. Carlini^j, J. Cha^c, R. Chrien^k, M. Christy^c, L. Cole^c, S. Danagouliau^d, D. Dehnhard^l, M. Elaasar^m, A. Empl^b, R. Ent^j, H. Fenker^j, Y. Fujii^a, M. Furic^f, L. Gan^c, K. Garrow^j, A. Gasparian^c, P. Gueye^c, M. Harvey^c, O. Hashimoto^a, W. Hinton^c, B. Hu^c, E. Hungerford^b, C. Jackson^c, K. Johnston^o, H. Juengst^l, C. Keppel^c, K. Lan^b, Y. Liang^c, V.P. Likhachev^p, J.H. Liu^l, D. Mack^j, A. Margaryan^h, P. Markowitz^q, J. Martoff^e, H. Mkrtchyan^h, S. N. Nakamura^a, T. Petkovic^f, J. Reinhold^q, J. Roche^r, Y. Sato^{a,c}, R. Sawafta^d, N. Simicevic^o, G. Smith^j, S. Stepanyan^h, V. Tadevosyan^h, T. Takahashi^a, K. Tanida^s, L. Tang^{c,j}, M. Ukai^a, A. Uzzle^c, W. Vulcan^j, S. Wells^o, S. Wood^j, G. Xu^b, H. Yamaguchi^a, C. Yan^j

(HNSS Collaboration)

^a*Tohoku University,*

Sendai, 980-8578, Japan;

^b*University of Houston,*

Houston, TX 77204;

^c*Hampton University,*

Hampton, VA 23668;

^d*North Carolina A&T State University,*

Greensboro, NC 27411;

^e*Temple University,*

Philadelphia, PA 19122;

^f*University of Zagreb, Zagreb, Croatia;*

^g*University of Bucharest,*

Bucharest, Romania;

^h*Yerevan Physics Institute,*

Yerevan, Armenia;

ⁱ*University of Maryland,*

College Park, MD 20742;

*^jThomas Jefferson National Accelerator Facility,
Newport News, VA 23606;*

*^kBrookhaven National Laboratory,
Upton, NY 11973;*

*^lUniversity of Minnesota,
Minneapolis, MN 55455;*

*^mSouthern University at New Orleans,
New Orleans, LA 70126;*

*ⁿRensselaer Polytechnic Institute,
Troy, NY 12180;*

*^oLouisiana Tech University,
Ruston, LA 71272;*

*^pUniversity of Sao Paulo,
Sao Paulo, Brazil;*

*^qFlorida International University,
Miami, FL 33199;*

*^rCollege of Williams and Mary,
Williamsburg, VA 23187;*

*^sUniversity of Tokyo,
Tokyo 113-0033, Japan*

(Dated: February 8, 2008)

Abstract

High energy CW electron beams at new accelerator facilities allow electromagnetic production and precision study of hypernuclear structure, and we report here on the first experiment demonstrating the usefulness of the $(e,e'K^+)$ reaction. This experiment is also the first to take advantage of the enhanced virtual photon flux available when electrons are scattered at approximately zero degrees. The observed resolution, ~ 900 keV, of the ${}^1_{{\Lambda}}\text{B}$ spectrum is the best yet attained using magnetic spectrometers. The positions of the major excitations are in agreement with theoretical predictions and the previous binding energy measurements.

PACS numbers:

One of the goals of hypernuclear spectroscopy has been to determine an effective Λ -Nucleon interaction[1] which is then tested against proposed Λ -Nucleon potentials folded into the nuclear many-body problem. A variety of spectroscopic data, particularly data sensitive to the effective spin-dependent potential parameters, is needed to help determine these parameters[2], and this will improve the theoretical treatment of the hadronic many-body system with strangeness and the knowledge of the free Λ -Nucleon interaction. The $(e,e'K^+)$ reaction has a large spin-flip component, producing states of non-natural parity, and substituting a Λ for a proton in the target nucleus. This produces hypernuclei charge symmetric to the previously employed (K^-, π^-) and (π^+, K^+) reactions.

The cross section for electroproduction can be written in a very intuitive form by separating out a factor, Γ , which multiplies the off-shell (virtual) photoproduction cross section. This factor may be interpreted as the virtual photon flux produced by (e,e') scattering[3, 4, 5, 6]. It has the form:

$$\Gamma = \frac{1}{2\pi} \left(\frac{\alpha}{-q_u^2} \right) \frac{E_\gamma}{1 - \epsilon} \left(\frac{E'_e}{E_e} \right). \quad (1)$$

In the above equation, ϵ is the polarization factor which vanishes for scattering at zero degrees, and $-q_u^2$ is the square of the 4-momentum of the virtual photon.

The virtual photoproduction cross section can then be approximated by the real photoproduction cross section at forward angles where the virtual photons are almost on the mass shell. The virtual photon flux factor, Γ , also has the feature that it is very forward peaked, so that for ~ 2 GeV electron energies, $\geq 50\%$ of the total flux is captured between scattering angles of 0 and 10mr[4, 5, 6]. Therefore if the electron spectrometer is placed at zero degrees, it needs only a small acceptance to assure that almost all the scattered electrons are transported to the focal plane, and that aberrations in the transport optics of the spectrometer system will be small and easily corrected. On the other hand, zero-degree-electroproduction uses essentially transverse photons, and it results in very high electron rates due to bremsstrahlung in the target[6].

The electron beam at Jlab is $\approx 60 \mu\text{m}$ in diameter with a momentum spread of $\Delta p/p \approx 10^{-5}$. At a beam energy of ~ 2 GeV, this introduces negligible error in the missing mass resolution. However, beam stability is a much more important issue, i.e. the requirement that whatever the beam momentum, it must be stable and reproducible for days (weeks)

during the course of an experimental run. In the experiment reported here, beam stability was accomplished to the level of a few parts in 10^{-4} by implementing a feed-back lock to the accelerator controls based on momentum measurements of the beam in the arcs of the accelerator. Changes in beam momentum were also recorded as each experimental run progressed, but corrections based on this information did not need to be applied.

The layout of the HNSS (E89-009) experiment is shown in Figure 1. An electron beam of primary energy ~ 1.8 GeV and $\sim 1\mu\text{A}$ current, strikes a thin target (20 mg/cm²) placed just before a small zero-degree dipole magnet. This magnet splits the scattered particles; the e^- of about 300 MeV/c into a split-pole spectrometer[7], ENGE, and the K^+ of about 1.2 GeV/c into the Short Orbit Spectrometer, SOS, which is fixed to the Hall C pivot at the Jefferson Laboratory. The splitting magnet is necessary in order to capture the scattered electrons and reaction kaons at very forward angles.




FIG. 1: The experimental plan view showing both the kaon spectrometer (SOS) and the electron spectrometer (ENGE)

The resolution for the experimental arrangement was expected to be 600-1000 keV. This was dominated by the SOS spectrometer, which was not designed for high resolution spectroscopy. However, the SOS was a short-orbit magnet (important as the decay length for a 1.2 GeV/c kaon was about 9m), was already mounted and tested at the Hall C pivot, and had the sophisticated particle identification[8], PID, package required to identify kaons within the expectedly large background of pions and positrons. Therefore, for the first experiment demonstrating the feasibility of this experimental technique, the SOS was chosen as the kaon spectrometer.

Two targets, 1) 22 mg/cm² ^{nat}C, and 2) 19 mg/cm² ⁷Li, were investigated, although a calibration target, 8.8 mg/cm² CH_x foil, was used to observe Λ and Σ production from hydrogen. Only the analysis of the C and CH_x data are reported here. The reaction of

interest, $(e,e'K^+)$, resulted in the production of the ${}^{12}_{\Lambda}\text{B}$ hypernucleus when a C nucleus was targeted. Beam intensities were tuned to produce an acceptable signal to accidental ratio, which for the C target was approximately $0.6\ \mu\text{A}$, or an experimental luminosity of $\sim 4 \times 10^{33}\ \text{cm}^{-2}\text{-s}^{-1}$. A beam current of $0.4\ \mu\text{A}$ was used with the CH_x target. To satisfy conflicting beam requirements when simultaneously operating several experiments in the different Halls, data had to be acquired at two different beam energies, 1721 and 1864 MeV. However, analysis has shown that within statistics, there are no differences in the resulting ${}^{12}_{\Lambda}\text{B}$ spectra.

Because of the high electron rates, the event trigger was the identification of a kaon in the SOS arm. The coincidence of this kaon with an electron in the ENGE spectrometer was completed in the off-line analysis. The number of positrons, pions, protons, and kaons per second from the C target detected in the SOS was 10^5 , 1.4×10^3 , 140, and 0.4 respectively. Therefore excellent particle identification was required, not only in the analysis, but also in the hardware trigger. The standard SOS detector package was used to identify kaons, and its description and operation have been previously discussed[8]. The large flux of positrons in the SOS came from the acceptance of scattering angles at 0° , where positrons from Dalitz pairs created in the target were observed. Positrons were easily identified and removed by the shower counter PID, but were used here to confirm the experimental resolution.

While the magnetic optics of the SOS were previously well studied, it was necessary to recalibrate this spectrometer because of the addition of the splitting magnet. For example, the angular acceptance of the SOS, normally 15 msr, was reduced to 6 msr by this magnet. The simulation of charged particle trajectories through this system of magnets and detectors used the program, RAYTRACE[9]. The parameters of the RAYTRACE code were determined by adjusting them so that the calculated multi-dimensional particle distributions matched those observed when the entrance angles and positions of reaction protons and pions from the target were restricted by a set of appropriately positioned holes in a tungsten plate (sieve slit) located between the splitting magnet and the SOS[10].

The electrons were detected near the focal plane of the Split-pole spectrometer by a set of silicon strip detectors (SSD) specifically designed for this experiment[11, 12]. Here only position along the focal plane was needed to obtain the required resolution. As it was also necessary to obtain a hit-position vs momentum calibration for the ENGE spectrometer, RAYTRACE parameters were adjusted to match the observed positions of the

$p(e, e'K^+)\Lambda$ or Σ peaks produced from the CH_x target. RAYTRACE also was used in a Monte Carlo program to determine the acceptance of the two spectrometers for the cross section calculation.

The calibration spectrum obtained from the CH_x target with the accidental background, as taken with the 1864 MeV beam, is shown in Figure 2. The Λ and Σ peaks are obvious, as well as an enhanced region just below the Λ peak. This enhancement is due to $(e, e'K^+)$ reactions on C in this target, and the threshold for ^{12}B production lies about 37.4 MeV below the Λ peak. The missing mass resolution in this spectrum, ~ 3.5 MeV (FWHM), is dominated by the kinematics due to the Λ recoil, and cannot be corrected because of the intrinsic angular resolution ($\sim 13\text{mr}$ FWHM) of the SOS spectrometer. This kinematic effect falls rapidly with target mass so that for the C target the contribution is small.




FIG. 2: The missing mass spectrum obtained from a CH_x target at an incident electron energy of 1864 MeV. The solid histogram is the accidental background as described in the text.

Although the Λ and Σ peaks in this spectrum were used to calibrate the focal plane of the SSD detector system, the known Λ photoproduction cross section could not be used for normalization because during irradiation the beam changed the hydrogen to carbon ratio of the CH_x target. The absolute value and shape of the accidental background was obtained by averaging over 8 accidental time peaks.

A measure of the experimental resolution can be obtained from the Dalitz pair data,

$A(e, e'[e^+e^-])A$, where the two electrons are observed in the ENGE and the positron in the SOS spectrometers, although this reaction did not fill the entire acceptance of the spectrometer. In this spectrum, the projected full energy peak has an experimental resolution of ~ 900 keV, which predicts a system resolution of ~ 890 keV after removing the resolution (calculated) of one electron by quadratures. This is consistent with the observed resolution in the hypernuclear spectrum.

The $^{12}\text{C}(e, e'K^+)_{\Lambda}^{12}\text{B}$ spectrum with the accidental background is shown in figure 3. Clearly evident are the ^{12}B hypernuclear excitations where a Λ is inserted in the s and p shells. The ground state doublet ($1^-; 2^-$) as well as the ($2^+; 3^+$) p-shell excitations are unresolved. There is also strength in the bound-state region between these excitations, although statistics are poor.




FIG. 3: The summed ^{12}B missing mass spectrum for both energies. The solid histogram is the measured accidental background, and the curve is a theoretical calculation, spread by 900 keV and overlaid on, not fit to, the data.

This spectrum is similar to that predicted by Motoba, *et al* [13, 14] and by Millener[15]. Reference [14] also calculates the excitation strengths for the photoproduction process of a kaon at an angle of 3 deg by a 1.3 GeV photon. The theoretical curve in the figure is generated by superimposing Gaussian peaks, having a FWHM of 900 keV below and 5 MeV above

15 MeV excitation energy, on a polynomial fit to the accidental background. The strength of the peaks is taken from reference [14] and their positions from [15]. The Millener[15] spectrum used an effective Λ -Nucleus interaction for the p-shell previously matched to other data, and thus should better represent the observed excitation energies. The curve in the figure is directly overlayed on (not fitted to) the data, without any shift in the energy scale.

The binding energy scale is determined from the position of the Λ and Σ peaks in the calibration spectrum. The $^{12}_{\Lambda}\text{B}$ binding energy is found to be 11.4 ± 0.5 MeV compared to the emulsion measurement of 11.37 MeV.

This spectrum can also be compared to the $\text{C}(\pi^+, \text{K}^+)_{\Lambda}^{12}\text{C}$ taken at KEK[16]. The resolution in this meson-induced spectrum is 1.45 MeV and shows the general shell structure as well as evidence for core excited states, albeit with better statistics but poorer resolution. Of course this reaction excites natural parity states, so although the overall shape of the spectrum is similar due to the small value of the spin-dependent ΛN effective interaction, the spin of the excited states differs. We note in particular, the strength between 5 to 8 MeV excitation which is observed in both data.

The differential cross section can be calculated as if it were photoproduction, by putting the virtual photons created in the (e, e') reaction on the mass shell. This in effect, averages the elementary (γ, K) reaction at ~ 1500 MeV over the approximately 100 MeV spread of virtual gamma energies. Since various factors entering the cross section calculation changed as a function of the incident energy, an independent analysis for each energy was required. The weighted average of these separate cross section measurements is, $140 \pm 17(\text{stat}) \pm 18(\text{sys})$ nb/sr, and is consistent with the value at both energies separately. It is also consistent with the theoretical prediction[14] of 152 nb/sr for the ground state.

The cross section normalization can be compared for consistency with two other experiments[17, 18]. Although these experiments did not resolve hypernuclear bound states, they did determine the effective number of protons participating in quasi-free production of Λ s from a C nucleus. To make this comparison, the background subtracted quasi-free missing mass spectrum of our data (not shown in the figure) was fitted by a polynomial function over a range of 23 MeV beginning at particle threshold. The ratio of the Σ to the Λ quasi-free production in this spectrum was assumed to be the same as that extracted from [17]. To compare to the previous experiments, the yield was corrected for acceptance, and differences in momentum transfer and energy[19]. We obtain an effective quasi-free cross

section of $4.55 \pm 0.19 \pm 0.46 \mu\text{b}/\text{sr}$ compared to $4.2 \pm 0.4 \mu\text{b}/\text{sr}$, [18]. From this cross section, we obtain 4.2 effective protons in quasi-free lambda production from C, in agreement with previous experiments and confirming our cross section normalization.

In summary, the first $(e,e'\text{K}^+)$ experiment has successfully observed the electroproduction of ${}^{12}_{\Lambda}\text{B}$. The missing mass resolution, ~ 900 keV, is consistent with the expected value, and is a factor of ~ 1.6 better than existing measurements. The spectrum shape and differential cross sections are also consistent with theory, and the binding energy is consistent with that previously measured by emulsion. Although there is strength in the core excited region, the statistical accuracy is not sufficient to extract detailed structure information. However the position and strength of the states are similar to those observed in the (π^+, K^+) reaction. A normalization based on the measured quasi-free production is also consistent with previous experiments[17, 18], and confirms our cross section value.

-
- [1] D. J. Millener, C. B. Dover, and A. Gal, Phys. Rev. **C28**, 2700(1988); T. Motoba, *et al*, Phys. Rev. **C38**, (1988)1322.
 - [2] D. J. Millener, Nucl. Phys. **A691**, 93c(2001).
 - [3] P. Barreau, *et al*, Nucl. Phys. **A402**, 515(1983).
 - [4] C. E. Hyde-Wright, W. Bertozzi, and J. M. Finn, Proc. 1985 CEBAF Summer Workshop, Newport News, VA, 1985.
 - [5] E. V. Hungerford, Prog. Theor. Phys. Supp., **117**, 135(1994).
 - [6] G. Xu and E. V. Hungerford, Preprint, University of Houston (2002).
 - [7] J. E. Spencer and H. A. Enge, Nucl. Inst. and Meth. **49**,181(1967).
 - [8] G. Niculescu, *et al*, Phys. Rev. Lett., **87**, 1805(1998).
 - [9] S. Kowalski and H. A. Enge, Nucl. Inst. and Meth. **A258**. (1987)407.
 - [10] L. Tang, C. Yan, E. V. Hungerford, Nucl. Inst. and Meth. **A366**, 259(1995).
 - [11] K. Lan, *et al*, IEEE Trans. Nucl. Sci. **46**, 1974(1999).
 - [12] T. Miyoshi, *et al*, to be published Nucl. Inst. and Meth.
 - [13] T. Motoba, M. Sotona, and K. Itonga, Prog. Theor. Phys. Supp., **117**, 123(1994).
 - [14] T. Motoba, Proceedings of the 8th Conference on Mesons and Light Nuclei, J. Adams, ed., AIP **603**, 125(2001); and T. Motoba, private communication.

- [15] D. J. Millener, Proc. of Jlab Workshop on Hypernuclear Physics with Electromagnetic Probes, Hampton University, Hampton, VA (1999).
- [16] T. Hasegawa, *et al*, Phys. Rev. Lett. **74**, 224(1995); H. Hotchi, *et al*, Phys. Rev. , **C64**, 044302(2001).
- [17] W. Hinton, PhD dissertation, Hampton University (1999).
- [18] H. Yamazaki, *et al*, Phys. Rev., **C52**, R1157(1999).
- [19] C. J. Bebek, *et al*, Phys. Rev. **D15**,594(1977).